

A DYNAMICAL MODEL FOR THE RESONANT MULTIPOLES AND THE Δ STRUCTURE

S. S. KAMALOV^a and Shin Nan YANG

Department of Physics, National Taiwan University, Taipei 10617, Taiwan

We show that recent experiment data for the ratios E_{1+}/M_{1+} and S_{1+}/M_{1+} can be explained in a dynamical model for electromagnetic production of pions, together with a simple scaling assumption for the bare $\gamma^*N\Delta$ form factors. Within our model we find that the bare Δ is almost spherical and the electric E2 and Coulomb C2 quadrupole excitations of the physical Δ are nearly saturated by pion cloud contribution in $Q^2 \leq 4.0 \text{ GeV}^2$.

1 Introduction

The study of excitations of the hadrons can shed light on the nonperturbative aspects of QCD. One case which has recently been under intensive study is the electromagnetic excitation of the $\Delta(1232)$ resonance. At low four-momentum transfer squared Q^2 , the interest is motivated by the possibility of observing a D state in the Δ . The existence of a D state in the Δ has the consequence that the Δ is deformed and the photon can excite a nucleon through electric $E2$ and Coulomb $C2$ quadrupole transitions. In a symmetric SU(6) quark model, the electromagnetic excitation of the Δ could proceed only via magnetic $M1$ transition. In pion electroproduction, $E2$ and $C2$ excitations would give rise to nonvanishing $E_{1+}^{(3/2)}$ and $S_{1+}^{(3/2)}$ multipole amplitudes. Recent experiments give nonvanishing ratio $R_{EM} = E_{1+}^{(3/2)}/M_{1+}^{(3/2)} \sim -0.03$ ¹ at $Q^2 = 0$ which has been widely taken as an indication of the Δ deformation.

At sufficiently large Q^2 , the perturbative QCD (pQCD) is expected to work. It predicts that only helicity-conserving amplitudes contribute at high Q^2 , leading to $R_{EM} = E_{1+}^{(3/2)}/M_{1+}^{(3/2)} \rightarrow 1$ and $R_{SM} = S_{1+}^{(3/2)}/M_{1+}^{(3/2)} \rightarrow \text{const.}$ This behavior in the perturbative domain is very different from that in the nonperturbative one. It is an intriguing question to find the region of Q^2 which signals the onset of the pQCD.

In a recent measurement², the electromagnetic excitation of the Δ was studied at $Q^2 = 2.8$ and 4.0 GeV^2 via reaction $p(e, e'p)\pi^0$. The extracted ratios R_{EM} and R_{SM} remain small and *negative*. This disagrees with the previous analysis³ of the earlier DESY data which gave small but *positive* R_{EM} and R_{SM} at $Q^2 = 3.2 \text{ GeV}^2$, though both analyses indicate that pQCD

^aPermanent address: Laboratory of Theoretical Physics, JINR Dubna, 141980 Moscow region, Russia

is still not applicable in this region of Q^2 . In this talk, we want to show⁴ that the recent data of Ref. 2 can be understood from the dominance of the pion cloud contribution at low Q^2 in both $E_{1+}^{(3/2)}$ and $S_{1+}^{(3/2)}$ multipoles, as predicted by a dynamical model^{5,6} for electromagnetic production of pion, together with a simple scaling assumption for the bare $\gamma^* N \Delta$ form factors.

2 Dynamical Model for $\gamma^* N \rightarrow \pi N$

The main feature of dynamical approach to pion photo- and electro-production^{5,6} is that the unitarity is built in by explicitly including the final state πN interaction in the theory, namely, t-matrix is expressed as

$$t_{\gamma\pi}(E) = v_{\gamma\pi} + v_{\gamma\pi} g_0(E) t_{\pi N}(E), \quad (1)$$

where $v_{\gamma\pi}$ is the transition potential operator for $\gamma^* N \rightarrow \pi N$, and $t_{\pi N}$ and g_0 denote the πN t-matrix and free propagator, respectively, with $E \equiv W$ the total energy in the CM frame.

Multipole decomposition of Eq. (1) gives the physical amplitude in channel α ⁶

$$t_{\gamma\pi}^{(\alpha)}(q_E, k; E + i\epsilon) = \exp(i\delta^{(\alpha)}) \cos \delta^{(\alpha)} \times \left[v_{\gamma\pi}^{(\alpha)}(q_E, k) + P \int_0^\infty dq' \frac{q'^2 R_{\pi N}^{(\alpha)}(q_E, q'; E) v_{\gamma\pi}^{(\alpha)}(q', k)}{E - E_{\pi N}(q')} \right], \quad (2)$$

where $\delta^{(\alpha)}$ and $R^{(\alpha)}$ are the πN scattering phase shift and reaction matrix in channel α , respectively; q_E is the pion on-shell momentum and $k = |\mathbf{k}|$ is the photon momentum.

The multipole amplitude in Eq. (2) manifestly satisfies the Watson theorem and shows that $\gamma\pi$ multipoles depend on the half-off-shell behavior of πN interaction. We remark that the use of K-matrix unitarization scheme as employed in, e.g., Ref. 7 would amount to approximating Eq. (2) with

$$t_{\gamma\pi}^{(\alpha)}(q_E, k; E + i\epsilon) = \exp(i\delta^{(\alpha)}) \cos \delta^{(\alpha)} v_{\gamma\pi}^{(\alpha)}(q_E, k). \quad (3)$$

The difference between Eqs. (2) and (3) lies in the fact that only the on-shell rescatterings are included in the K-matrix unitarization scheme.

In the resonant (3,3) channel in which $\Delta(1232)$ plays a dominant role, the transition potential $v_{\gamma\pi}$ consists of two terms

$$v_{\gamma\pi}(E) = v_{\gamma\pi}^B + v_{\gamma\pi}^\Delta(E), \quad (4)$$

where $v_{\gamma\pi}^B$ is the background transition potential which includes Born terms and vector mesons exchange contributions, as described in Ref. 8. The second term of Eq. (4) corresponds to the contribution of bare Δ .

With Eq. (4), we may decompose Eq. (1) in the following way

$$t_{\gamma\pi}(E) = t_{\gamma\pi}^B + t_{\gamma\pi}^\Delta(E), \quad (5)$$

where

$$t_{\gamma\pi}^B(E) = v_{\gamma\pi}^B + v_{\gamma\pi}^B g_0(E) t_{\pi N}(E), \quad (6)$$

$$t_{\gamma\pi}^\Delta(E) = v_{\gamma\pi}^\Delta + v_{\gamma\pi}^\Delta g_0(E) t_{\pi N}(E). \quad (7)$$

Here $t_{\gamma\pi}^B$ includes contributions from the nonresonant background and renormalization on the vertex $\gamma^* N \Delta$ due to πN scattering. The advantage of such a decomposition is that all the processes which start with the electromagnetic excitation of the bare Δ are summed up in $t_{\gamma\pi}^\Delta$.

Multipole decomposition of Eq. (6) takes the same form as Eq. (2) and is used to calculate the multipole amplitudes $M_{1+}^B(W, Q^2)$, $E_{1+}^B(W, Q^2)$ and $S_{1+}^B(W, Q^2)$ with $R_{\pi N}^{(\alpha)}(q_E, q'; E)$ obtained from a meson exchange model⁹ for πN interaction. Note that to make the principal value integration in Eq. (2) associated with $v_{\gamma\pi}^B$ convergent, we introduce an off-shell dipole form factor with cut-off parameter $\Lambda=440$ MeV. The gauge invariance, violated due to the off-shell rescattering effects, is restored by the substitution $J_\mu^B \rightarrow J_\mu^B - k_\mu k \cdot J^B / k^2$, where J_μ^B is the electromagnetic current corresponding to the background contribution $v_{\gamma\pi}^B$.

3 $\gamma^* N \leftrightarrow \Delta$ transition form factors

Now let us consider the Δ resonance contribution $t_{\gamma\pi}^\Delta$ in Eq. (5). In keeping with the standard way of experimental analysis and constituent quark model (CQM) calculations, we describe the resonant multipoles $t_{\gamma\pi}^{\Delta, \alpha}$ with a Breit-Wigner type of energy dependence, as was done in the isobar model of Ref. 8,

$$t_{\gamma\pi}^{\Delta, \alpha}(W, Q^2) = \bar{\mathcal{A}}_\alpha^\Delta(Q^2) \frac{f_{\gamma\Delta} \Gamma_\Delta M_\Delta f_{\pi\Delta}}{M_\Delta^2 - W^2 - i M_\Delta \Gamma_\Delta} e^{i\phi}, \quad (8)$$

where $f_{\pi\Delta}(W)$ is the usual Breit-Wigner factor describing the decay of the Δ resonance with total width $\Gamma_\Delta(W)$ and physical mass $M_\Delta=1232$ MeV. The expressions for $f_{\gamma\Delta}$, $f_{\pi\Delta}$ and Γ_Δ are taken from Ref. 8. The phase $\phi(W)$ in Eq. (8) is to adjust the phase of $t_{\gamma\pi}^{\Delta, \alpha}$ to be equal to the corresponding pion-nucleon scattering phase $\delta^{(33)}$. Note that at the resonance energy $\phi(M_\Delta) = 0$.




Figure 1: Real and imaginary parts of the, **(a)** $M_{1+}^{(3/2)}$, and **(b)** $E_{1+}^{(3/2)}$ multipoles. The open and full circles are the results from the Mainz dispersion relation analysis¹⁰ and from the VPI analysis¹¹, respectively. Notations for the curves are given in the text.

The main parameters in the bare $\gamma^* N \Delta$ vertex are the $\bar{\mathcal{A}}^\Delta(Q^2)$'s in Eq. (8). For the magnetic dipole $\bar{\mathcal{M}}^\Delta$ and electric quadrupole $\bar{\mathcal{E}}^\Delta$ transitions, they are related to the conventional electromagnetic helicity amplitudes $A_{1/2}^\Delta$ and $A_{3/2}^\Delta$ by

$$\bar{\mathcal{M}}^\Delta(Q^2) = -\frac{1}{2}(A_{1/2}^\Delta + \sqrt{3}A_{3/2}^\Delta), \quad \bar{\mathcal{E}}^\Delta(Q^2) = \frac{1}{2}(-A_{1/2}^\Delta + \frac{1}{\sqrt{3}}A_{3/2}^\Delta). \quad (9)$$

At the photon point $Q^2 = 0$, the bare amplitudes $\bar{\mathcal{M}}^\Delta(0)$ and $\bar{\mathcal{E}}^\Delta(0)$ of Eq. (8) are determined from the best fit to the results of the recent analyses of Mainz¹⁰ (open circles) and VPI group¹¹ (full circles), as shown in Fig. 1. The dashed curves denote the contribution from $t_{\gamma\pi}^B$ only. The dotted curves correspond to the K-matrix approximation to $t_{\gamma\pi}^B$, namely, without the inclusion of principal value integral term. Solid curves are the full results of our calculation with bare Δ excitation.

The numerical values for $\bar{\mathcal{M}}^\Delta$ and $\bar{\mathcal{E}}^\Delta$ and the helicity amplitudes, at $Q^2 = 0$, are given in Table 1. Here we also give "dressed" values obtained using K-matrix approximation for $t_{\gamma\pi}^B$. One notices that the bare values determined above for the helicity amplitudes amount to only about 60% of the corresponding dressed values and are close to the predictions of the CQM, as pointed out by Sato and Lee¹². The large reduction of the helicity amplitudes from the dressed to the bares ones results from the fact that the principal value

integral part of $t_{\gamma\pi}^B$, which represents the effects of the off-shell pion rescattering, contributes approximately half of the M_{1+} as indicated by the dashed curves in Fig. 1.

Table 1: Comparison of the "bare" and "dressed" values for the amplitudes $\bar{\mathcal{A}}^\Delta$, $A_{1/2}^\Delta$ and $A_{3/2}^\Delta$ (in $10^{-3} \text{ GeV}^{-1/2}$).

Amplitudes	"bare"	"dressed"	PDG
$\mathcal{M}^\Delta$	158 ± 2	289 ± 2	293 ± 8
$\bar{\mathcal{E}}^\Delta$	0.4 ± 0.3	-7 ± 0.4	-4.5 ± 4.2
$A_{1/2}^\Delta$	-80 ± 2	-134 ± 2	-140 ± 5
$A_{3/2}^\Delta$	-136 ± 3	-256 ± 2	-258 ± 6

We now turn to the Q^2 evolution of the multipoles $\bar{\mathcal{M}}^\Delta(Q^2)$ and $\bar{\mathcal{E}}^\Delta(Q^2)$. In the present work, we parametrize the Q^2 dependence of the dominant $\bar{\mathcal{M}}^\Delta$ amplitude by

$$\bar{\mathcal{M}}^\Delta(Q^2) = \bar{\mathcal{M}}(0) \frac{|\mathbf{k}|}{k_\Delta} (1 + \beta Q^2) e^{-\gamma Q^2} G_D(Q^2), \quad (10)$$

where G_D is the nucleon dipole form factor. For the small $\bar{\mathcal{E}}^\Delta$ and $\bar{\mathcal{S}}^\Delta$ amplitudes, we follow Ref. 8 and assume that they have the same Q^2 dependence as $\bar{\mathcal{M}}^\Delta$ (*scaling assumption*). This is motivated by the scaling law which has been observed for the nucleon form factors.

We remind the reader that, in contrast to Ref. 8, amplitudes $\bar{\mathcal{M}}_{1+}$ and $\bar{\mathcal{E}}_{1+}$ and the corresponding helicity amplitudes in Eq. (9) correspond to the "bare" $\gamma N \Delta$ transition. For the real photon, they are equal to the standard $M1$ and $E2$ amplitudes of the $\Delta \leftrightarrow N\gamma$ transition defined in accordance with the convention of the Particle Data Groups. At the resonance energy, they can be easily expressed in terms of the Dirac-type form factors g_1 and g_2 used in Ref. 7, or Sachs-type form factors G_M and G_E used in Ref. 12. The relation between $\bar{\mathcal{M}}_{1+}$ amplitude and the bare G_M form factors is as follows

$$\bar{\mathcal{M}}_{1+}(Q^2) = \frac{e}{2m} \frac{|\mathbf{k}|}{k_\Delta} \sqrt{\frac{k_\Delta M_\Delta}{m}} G_M(Q^2), \quad (11)$$

where $k_\Delta = (M_\Delta^2 - m^2)/2M_\Delta$ with m and M_Δ denoting the nucleon and Δ mass, respectively. Expression for the electric amplitude is similar, but with opposite sign. Relation between physical $M_{1+}^{(3/2)}$ multipole and experimentally measured G_M^* form factor is given by Eq. (24) of Ref. 8. Note that we employ the "Ash" definition¹⁴ for the G_M^* which differs from the Δ form factor used in Ref. 2 by a factor $f = \sqrt{1 + Q^2/(m + M_\Delta)^2}$, i.e., $G_M^*(our) = G_M^*(Ref. 2)/f$.

4 Results and Discussion

Using the β and γ in Eq. (10) as free parameters, we fit the recent experimental data² as well as old one quoted in Ref. 8 on the Q^2 dependence of the $M_{1+}^{(3/2)}$ or equivalently, the G_M^* form factor. Our result is shown in Fig. 2. The obtained values for the β and γ parameters are: $\beta = 0.44 \text{ GeV}^{-2}$ and $\gamma = 0.38 \text{ GeV}^{-2}$. Here the dashed curve corresponds to contribution from the bare Δ , i.e., $t_{\gamma\pi}^\Delta$ of Eq. (7). The results for the ratios $R_{EM} = E_{1+}^{(3/2)}/M_{1+}^{(3/2)}$




Figure 2: The Q^2 dependence of (a) $Im M_{1+}^{(3/2)}$ with corresponding G_M^* form factor, and (b) ratios $E_{1+}^{(3/2)}/M_{1+}^{(3/2)}$ and $S_{1+}^{(3/2)}/M_{1+}^{(3/2)}$ at $W=1232$ MeV. The full and dashed curves are the results for the "dressed" and "bare" $\gamma N\Delta$ vertexes, respectively. Experimental data as quoted in Ref. 8. The new data at $Q^2=2.8$ and 4.0 (GeV/c)² are from Ref. 2.

and $R_{SM} = S_{1+}^{(3/2)}/M_{1+}^{(3/2)}$ are shown in the right column of Fig. 2. It is seen that they are in good agreement with the results of the model independent analysis of Ref. 2 up to Q^2 as high as 4.0 GeV^2 . Note that since the bare values for the E2 and C2 excitations are small, the absolute values and shape of these ratios are determined, to a large extent, by the pion rescattering contribution. The bare Δ excitation contributes mostly to the $M_{1+}^{(3/2)}$ multipole.

Pion cloud has been found to play an important role in hadron structure in many studies. For example, in the cloudy bag model (CBM), a reasonably good agreement^{15,16} with the measured R_{EM} can be obtained with bag radius of $R = 0.6 - 0.8 \text{ fm}$. In a recent improved CBM calculation¹⁷, where the relativistic effects and CM motion were better treated, it was also found that the pion cloud gives large contribution to $G_M^*(Q^2)$. It is generally concluded

that the smaller the bag radius, the larger the pion cloud contribution. Similar conclusion is reached with respect to the proton EM form factors within the cloudy bag model. In fact, it has been found¹⁸ that the recent Jlab data¹⁹ on the scaling violation in the electromagnetic form factors can be explained within CBM with $R = 0.7 \text{ fm}$. One might then be tempted to interpret our result as another indication of preferring a smaller bag radius. However, at these small bag radii, the pion field is so strong that the use of the perturbative approach employed in these CBM studies is questionable. In this connection, we should mention that in a nonperturbative calculation within a chiral chromodielectric model and a linear σ -model, Fiolhais, Golli, and Sirca²⁰ reached a similar conclusion as ours, namely, the large experimental values of R_{EM} and R_{SM} could be explained in terms of the pion contribution alone.

In nonrelativistic CQM it is well known that, the values of R_{EM} and R_{SM} obtained with a D -state admixture in the Δ generated by one-gluon-exchange hyperfine interaction are in general too small. It has recently been suggested by the Tübingen group²¹ that this problem can be fixed with the inclusion of exchange currents in the calculation. The reason is that two-body exchange currents could flip the spins of the two quarks in the nucleon to convert it into a Δ . Such a transition would just be a transition between the S-states in the nucleon and Δ , and accordingly is greatly enhanced. We would like to point out here that the diagram involving a two-body exchange current induced by the exchange of a pion between two quarks in the nucleon can also be interpreted as pion rescattering effects, as considered in the current study.

5 Summary

In summary, we calculate the Q^2 dependence of the ratios E_{1+}/M_{1+} and S_{1+}/M_{1+} in the $\gamma^*N \rightarrow \Delta$ transition, with the use of a dynamical model for electromagnetic production of pions. We find⁴ that both ratios E_{1+}/M_{1+} and S_{1+}/M_{1+} remain small and negative for $Q^2 \leq 4.0 \text{ GeV}^2$. Our results agree well with the recent measurement of Frolov et al.², but deviate strongly from the predictions of pQCD. Our results indicate that the bare Δ is almost spherical and hence very difficult to be directly excited via electric E2 and Coulomb C2 quadrupole excitations. The experimentally observed $E_{1+}^{(3/2)}$ and $S_{1+}^{(3/2)}$ multipoles are, to a very large extent, saturated by the contribution from pion cloud, i.e., pion rescattering effects. It remains an intriguing question, both experimentally and theoretically, to find the region of Q^2 which will signal the onset of pQCD.

Acknowledgments

This work is supported in part by the NSC/ROC under the grant no. NSC 88-2112-M002-015.

References

1. R. Beck et al., *Phys. Rev. Lett.* **78**, 606 (1997); G. Blanpied et al. *Phys. Rev. Lett.* **79**, 4337 (1997).
2. V.V. Frolov et al., *Phys. Rev. Lett.* **82**, 45 (1999).
3. V. Burkert and L. Elouadrhiri, *Phys. Rev. Lett.* **75**, 3614 (1995).
4. S.S. Kamalov and S.N. Yang, *Phys. Rev. Lett.* **83**, 4494 (1999).
5. H. Tanabe and K. Ohta, *Phys. Rev. C* **31**, 1876 (1985).
6. S.N. Yang, *J. Phys. G* **11**, L205 (1985).
7. R. Davidson, N. C. Mukhopadhyay and R. Wittman, *Phys. Rev.* **D43**, 71 (1991).
8. D. Drechsel, O. Hanstein, S.S. Kamalov and L. Tiator, *Nucl. Phys. A* **645**, 145 (1999).
9. C.T. Hung, S.N. Yang and T.-S.H. Lee, *J. Phys. G* **20**, 1531 (1994); C. Lee, S.N. Yang and T.-S.H. Lee, *ibid.* *G* **17**, L131 (1991).
10. O. Hanstein, D. Drechsel, and L. Tiator, *Nucl. Phys. A* **632**, 561 (1998).
11. R.A. Arndt, I.I. Strakovsky and R.L. Workman, *Phys. Rev. C* **53**, 430 (1996) (SP97 solution of the VPI analysis).
12. T. Sato and T.-S.H. Lee, *Phys. Rev. C* **54**, 2660 (1996); T.-S. H. Lee, in *N* Physics*, eds. T.-S.H. Lee and W. Roberts, p. 19, (World Scientific, Singapore 1997).
13. R. Davidson, N. C. Mukhopadhyay and R. Wittman, *Phys. Rev. Lett.* **56**, 804 (1986).
14. W. W. Ash, *Phys.Lett.* **24B**, 165 (1967).
15. G. Kälbermann and J.M. Eisenberg, *Phys. Rev. D* **28**,71 (1983).
16. K. Bermuth, D. Drechsel, L. Tiator, and J.B. Seaborn, *Phys. Rev. D* **37**, 89 (1988).
17. D.H. Lu, A.W. Thomas, and A.G. Thomas, *Phys. Rev. C* **55**, 3108 (1997).
18. D.H. Lu, S.N. Yang, and A.W. Thomas, *J. Phys. G* (2000), in print.
19. M. K. Jones et al, *Phys. Rev. Lett.* **84** 1398 (2000).
20. M. Fiolhais, B. Golli, and S. Sirca, *Phys. Lett. B* **373** 229 (2000).
21. A.J. Buchmann, E. Hernández, U. Meyer, and A. Faessler, *Phys. Rev. C* **58**, 2478 (1998); A.J. Buchmann, "Baryons' 98", eds. D. W. Menze and B. Ch. Metsch, World Scientific, Singapore, 1999, p. 731 (hep-ph/9909385); and this proceeding.